

This Free E-Book is brought to you by Natural-Aging.com.

100% Effective Natural Hormone Treatment
Menopause, Andropause And Other Hormone Imbalances
Impair Healthy Healing In People Over The Age Of 30!

Breast Cancer Statistics - How Breast Cancer Survival Rates Increased 50%

By Olinda Rola

Breast cancer statistics show that over 1.2 million persons will be diagnosed with breast cancer

worldwide this year, according to the World Health Organization. For breast cancer and prevention, it has long been known that regular physical activity has been shown to decrease the likelihood of having breast cancer. What has not been known or studied has been the effect of regular physical activity on the breast cancer survival rates or likelihood of death in women that already have breast cancer. That is, until now.

The breast cancer statistics and findings as reported by the American Medical Associations Journal of the American Medical Association (JAMA) in May 2005 were astounding! Certain participants in the study of women with Stage I, II or III breast cancer achieved a 50% reduction in the death rate from breast cancer.

Here are these breast cancer statistics: the journal reported that in the study 2,987 female registered nurses had been diagnosed with breast cancer during the years 1984–1998. What the study found was that the women who had physical activity equivalent to walking at a steady pace of 2.0–2.9 miles per hour for 3–5 hours a week had a death rate of only 50% of the death rate of women who had physical activity equivalent to walking less than one hour a week. The conclusion of the breast cancer statistics in the study was that physical activity after breast cancer has been diagnosed may reduce the risk of death from breast cancer. The study found that there was little evidence of any relation between increased physical activity and increased benefit.

It's time to dust off those walking shoes!

As a physical activity, walking can be done almost anytime by anyone anywhere. All that is needed is a good pair of walking shoes. Walking is fun and reduces stress. As for injuries, walking has the lowest injury rate of all the various kinds of exercise.

You can walk with a partner, friend, family member or dog, maybe even a neighbors dog. Or you can walk with your favorite headset and music. If you are walking outdoors with a headset, keep one ear open to hear the sounds around you.

Breast Cancer Statistics – How Breast Cancer Survival Rates Increased 50%

As for basic walking tips:

- As you begin regular walking, take it easy. Standard advice is to check with your physician before starting any exercise program. If it has been years or decades since you walked regularly, perhaps you can begin with 5 minutes of walking and slowly increase your time and distance.
- Walking at a pace of 75-95 steps a minute will have you walking at a speed of about 2–3 miles per hour.
- Walk with your head up, looking out in front of you. Do not walk looking down right in front of you except to navigate any obstacles.
- Really take it easy the first 5 minutes of walking to warm up. Afterwards, gently stretch for 5-10

minutes while your muscles are warm.

- Practice good walking form. Your arms should swing naturally in the direction you are walking, not from side to side across your body. Your foot should strike the ground on your heel, then a rolling motion forward toward the ball of your foot, then pushing off with your toes.

And here are some basic walking shoes tips:

- Buy your walking shoes from a sporting shoes store with large selections. Doing so will give you plenty of choices. And buy your walking shoes later in the day when your feet will be larger.
- Buy cushioned, supportive walking shoes. To see if a shoe is supportive, do this test: take a shoe and turn it upside down. Holding each end of the shoe, try to fold it. If you find the shoe bends in the middle, then that shoe is not a supportive shoe. A supportive shoe should bend where your foot normally bends, near your toes.
- You should allow the width of your index finger between the end of your shoe and the end of your longest toe, or about one–half inch.
- Buy two pairs of walking shoes, one for home and one for the car or workplace. And if one pair gets wet, you can use the other pair that day.

Walking is the closest thing to the perfect exercise. In today's fast–paced society, regular walking can be a welcomed break from the stress of the day. Maybe you will get to know your neighborhood or neighbors better. There may be walking trails you have never seen but wanted to.

Wherever and however you choose to walk, not only can the experience be fun, you will know you are being good to your body in a variety of ways. Besides the incredible breast cancer statistics and findings of the breast cancer study, walking helps with weight control and bone strength, elevates mood, helps build and maintain healthy muscles, joints and heart. With so many great health benefits, why not get started walking today!

Copyright 2005 InfoSearch Publishing

Olinda Rola is President of InfoSearch Publishing and webmaster of

<http://www.safemenopausesolutions.com>

– a website of natural treatments for a variety of health

problems. For information about preventing breast cancer, breast cancer treatment and related articles, visit the website to read more.

Reduce Breast Cancer By 45%!

By Camry James

A US study of 13,000 patients showed Tamoxifen reduced the rate of expected cancers from one in 130 to one in 236 – a cut of about 45%. The American researchers ended their trial early when they said the drug's benefits became overwhelmingly obvious.

Other studies found that tamoxifen AND chemotherapy improved survival improved survival rates by about 40-50% compared to taking one treatment or the other.

Should women taking tamoxifen avoid pregnancy?

Yes. Tamoxifen may make premenopausal women more fertile, but doctors advise women on tamoxifen to avoid pregnancy because animal studies have suggested that the use of tamoxifen in pregnancy can cause fetal harm. Women who have questions about fertility, birth control, or pregnancy should discuss their concerns with their doctor.

Does tamoxifen cause a woman to begin menopause?

Tamoxifen does not cause a woman to begin menopause, although it can cause some symptoms that are similar to those that may occur during menopause. In most premenopausal women taking tamoxifen, the ovaries continue to act normally and produce estrogen in the same or slightly increased amounts.

Do the benefits of tamoxifen in treating breast cancer outweigh its risks?

The benefits of tamoxifen as a treatment for breast cancer are firmly established and far outweigh the potential risks. Patients who are concerned about the risks and benefits of tamoxifen or any other medications are encouraged to discuss these concerns with their doctor.

Tamoxifen (ta-MOX-i-fen) is a medicine that blocks the effects of the estrogen hormone in the body. It is used to treat breast cancer in women or men. It may also be used to treat other kinds of cancer, as determined by your doctor.

Breast Cancer Statistics – How Breast Cancer Survival Rates Increased 50%

Tamoxifen also may be used to reduce the risk of developing breast cancer in women who have a high risk of developing breast cancer. Women at high risk for developing breast cancer are at least 35 years of age and have a combination of risk factors that make their chance of developing breast cancer 1.67% or more over the next 5 years. Your doctor will help to determine your risk of developing breast cancer.

Tamoxifen has been proven to be the cure to Breast Cancer. We urge you to find out more about Tamoxifen at

<http://Tamoxifen.eask.info>



This Free E-Book has been brought to you by Natural-Aging.com.

**[100% Effective Natural Hormone Treatment](#)
Menopause, Andropause And Other Hormone Imbalances
Impair Healthy Healing In People Over The Age Of 30!**